

UDK 630*52:311.2(497.6)
630*52:007.5(497.6)

WEIGHTED FUNCTIONS in the k -NN ESTIMATES of GROWING STOCK in HIGH FOREST in BOSNIA

Ponderske funkcije u k -nn procjenama drvne zalihe u visokim šumama Bosne

**Azra Čabaravdić¹, Dieter R. Pelz², Gherardo Chirici³, Christian Kutzer⁴,
Ernada Ćatić⁵, Hamid Delić⁵**

Abstract

Last decades permanent researches clarify possibilities for forest resource estimation based on terrestrial measurement and remote sensing. The most often the non-parametrical k -NN method is used integrating local estimates from terrestrial measurement and spectral Landsat data. In this paper the weighting functions of the k -NN related to value differences and distances were examined in a case of high forest in site Konjuh in Bosnia. It is found that weighting Euclidean distance has not resulted with efficiency increase. Procentual RMSE's of growing stock showed higher values for weighted estimates on the pixel level. Classified volume estimates on aggregated level compared with volumes from intensive regular forest inventory achieved moderate level of agreement. The agreements between volume estimates are almost perfect regardless on weighting functions. Obtained results point out unweighted estimates as reported in several cases.

Key words: *growing stock, Landsat TM+, k -NN estimates, Euclidian distance, weighting function*

¹ Faculty of Forestry, University of Sarajevo, Zagrebačka 20, 71000 Sarajevo, Bosnia-Herzegovina

² Department for Forest Biometry, Faculty of Forest and Environmental Sciences, Albert-Ludwigs-University Freiburg, Tennenbacher Straße 4 79085 Freiburg i. Br., Germany

³ Dipartimento di Scienze e Tecnologie per l'Ambiente e il Territorio, Università degli Studi del Molise, C.da Fonte Lappone - 86090 Pesche (IS), Italy

⁴ Ph.D. (Faculty of Forest and Environmental Sciences, Albert-Ludwigs-University Freiburg, Germany)

⁵ M.Sc. (Faculty of Forestry, University of Sarajevo, Zagrebačka 20, 71000 Sarajevo, Bosnia-Herzegovina)

INTRODUCTION - Uvod

Multisource forest inventory on the global, regional, local and small scale level become the imperative in modern forest governance and management. Recent possibilities in forest resource assessment resulted with estimations and classifications based on all available information and data. Sampling methods, field measurements, remote sensing, geoinformation system have been examined permanently. Their possibilities are reflected in forest resource estimations and mapping on large scale (GALLAUN ET AL., 2010; NABUURS, 2010).

Number of studies confirmed acceptable results based on non-parametric estimation method (*k-NN*) for forest inventory on large area (FRANCO-LOPEZ ET AL. 2001, McROBERTS ET AL. 2002, HAPPANEN ET AL. 2004, NILSSON 1997, REESE ET AL., 2003). The *k-NN* estimation was introduced by KILKKI AND PAIVINEN (1987) and adapted for Finnish national forest inventory (MS-NFI) since 1990. The MS-NFI was assigned as image-aided multisource method. During last the inventories the method was enhanced improving *k-NN* estimation procedures aiming to deliver as large so small area estimates and maps (TOMPPPO ET AL., 2001; GJERTSEN&ERIKSEN, 2004; KOUKAL, 2004; KANGAS&MALTAMO, 2006; MASELLI ET AL., 2005; FINLEY ET AL., 2006; FINLEY ET AL., 2008; BAFFETTA ET AL., 2009, 2011). The method is suitable for carbon cyclus and other applied research in forestry and environmental sciences (KÖHL ET AL., 2008; STÜMER ET AL., 2010; HÄRKÖNEN ET AL., 2011).

Basic *k-NN* estimation is based on a heuristic model and the estimation is not design-based. The estimates were based on local estimations of forest attributes and corresponding pixel values in multidimensional space. Firstly the reference sample plot method was developed where the term “k” was one. In the application it was not possible to control weight of the individual plot (at the population level) in the estimation procedure (KANGAS&MALTAMO, 2006). Several papers reported about improvements of methodological approach related to weighting functions. Firstly, the *k-NN* estimates on the pixel level were based on unweighted and inverse distance-weighted equations. Then optimization method based on a genetic algorithm was used in order to find the variable weights. The method was assigned as *ik-NN* (improved *k-NN* method) (TOMPPPO&HALME, 2004). The alternative to the *k-NN* methods have been developed and examined (Stümer& 2010). Recent advances and emerging issues in nearest neighbor’s techniques are reviewed for four topic areas: (1) distance metrics, (2) optimization, (3) diagnostic tools, and (4) inference (McROBERTS, 2011).

Past studies in Bosnia resulted with acceptable *k-NN* estimates for growing stock for some planning unit levels (the high forest with natural regeneration, the narrow forest categories and some management classes) (ČABARAVDIĆ, 2007). Here were compared *k-NN* estimates based on weighted Euclidean, Mahalanobis and modified Mahalanobis distance (MD) in spectral space. Further research questions could be related to the growing stock estimation considering possibilities of applying different weighting functions on Euclidian distance numerically or by some more advanced procedures. The present paper deals with the application of weighting functions related to

Euclidian distance and value difference in the non-parametrical k -NN method in a test forest inventory carried out in the Konjuh Kladanj area in B&H. The final aim is to set up an optimal k -NN model based on weighted Euclidian distance to spatialize the growing stock on the basis of Landsat 7 ETM+ imagery and to obtain if the application of weighting functions would increase estimation efficiency.

MATERIALS AND RESEARCH METHODS – Materijal i metod

The test site for this research is located in the nord-east Bosnia representing high forest with natural regeneration.

The total land area of Bosnia and Herzegovina (B&H) is approximately 51,500 square kilometers, of which about half is forested. Broadleaf trees dominate two-third of the forested landscape. Sixty percent is high forest, mainly mixed broadleaf-conifer stands or pure stands of oaks, beech, spruce, pines or fir. In the Federation of Bosnia and Herzegovina, forested area is 1,135,931 ha, with 56.3% of high forests with natural regeneration. Mixed beech, fir and spruce forests in central Bosnia are assigned as forests of the highest productivity and high structural and ecological stability. For the present research was chosen a test area defined as an administrative forest management unit named Forest Economic Management Unit “Konjuh” Kladanj with about 26,000 ha of forests. The high forest with natural regeneration dominates covering about 19,000 ha.




Figure 1. Test area Konjuh Kladanj
Slika 1. Područje istraživanja Konjuh Kladanj

For the purposes of the scientific research a pilot forest inventory was made in the test area. Ground sampling was configured on the basis of a single-stage systematic cluster sampling design. Based on UTM georeferenced reference system, a 2 kilometric sample grid was generated to locate clusters (Figure 2).




Figure 2. Sampling plan
Slika 2. Plan uzorka

Plots of 25 m in radius were aggregated into square-shaped clusters (Figure 3). On the grid clusters made of 8-sample plots distributed in square were displaced at a distance of 400 m one from the others.

A first selection of forest clusters was carried out on the basis mine-free land. After this selection, 3 clusters were classified as non-accessible and excluded from further analysis.

A total number of 253 sample plots remained, 235 were selected for the experimental test because belonging to the high forest with natural regeneration. The field work for acquiring forest inventory data was conducted in the summer 2003. Registered information and data were used to describe sampling plots and to determine local estimates of the most important forest attributes (growing stock, number stem and growth) and their confidence intervals. Satellite data were from Landsat 7 ETM+ 187-029 scene acquired in the august 2000 with 7 multispectral bands of 30 m of geometric resolution. Here are used 1st-5th and 7th spectral channels. In order to acquire information each of the 235 sampling units was inventoried in the field. Then spectral values of the corresponding pixels were extracted and recorded.

The *k*-NN method

The *k*-Nearest Neighbors (*k*-NN) classifier is one of the most widely known and used non-parametric classification procedures introducing in combined forest inventories by KATILA&TOMPPPO (2001) and FRANCO-LOPEZ ET AL. (2001). Later the method is investigated intensively and adjusted in many studies confirming its applicability in estimation procedures related to compiled forest and spectral data.

Detailed description of the method used in this paper is given in STÜMER (2004) and KUTZER (2008): “With the k-NN method are obtained forest attributes estimates calculating the average value of the k neighbour samples. The neighbours are weighted by the distance value, which describes the spectral similarity. Each pixel contains spectral information for each channel in the form of a digital value. The spectral difference between two pixels is defined with the use of a metrics. A common distance value is the Euclidean distance, $d_{(i)p}$, which has to be calculated from the target pixel p, to every sample pixel i, for which a terrestrial observation is available. If x_1 and x_2 are the characteristic vectors of two pixels whose similarity have to be tested, the Euclidean distance $d(x_1, x_2)$ between them is

$$d'_{(x_1, x_2)} = \left[\sum_{j=1}^N (x_{1j} - x_{2j})^2 \right]^{1/2}$$

where N is the number of the spectral components (e.g. used channels). A generalized Euclidean distance is described as the Minkowski-r-distance (BORTZ, 1993). By replacing the exponent 2 (respectively $1/2$) with r (respectively $1/r$), a generalization of the Euclidean distance equation (5.1) is the result:

$$d'_{(x_1, x_2)} = \left[\sum_{j=1}^N (x_{1j} - x_{2j})^r \right]^{1/r}$$

From $r = 1$, the City-Block-Metric results (BORTZ, 1993), of which the distance of two points results from the sum of the attribute differences.

By variation of the metric coefficient, attribute differences become weighted differently. With $r = 1$, all attribute differences are weighted equally, irrespectively of their term. With $r = 2$, bigger differences get a stronger loading compared to smaller differences and so on.

Also, the variability of the spectral information varies within the channels. To consider channels with a wide variability in the reflex reflection and to weight their influence on the differentiation of attribute classes of an attribute, the parameter a_j for the weighting of the channels has been introduced (FRANCO-LOPEZ ET AL., 2001):

$$d_{(x_1, x_2)} = \left[\sum_{j=1}^N a_j^r (x_{1j} - x_{2j})^r \right]^{1/r}$$

If the parameter a_j for $j = 1, \dots, N$ is chosen equal to 1, all channels have the same weight when calculating the distance. However, each channel can adequately be linked by a weighting a_j .

The $k = 1$ to $k = n$ nearest spectral neighbours, i.e. pixel with corresponding terrestrial observations are used for the following analyses.

$$d_{(1),p} \leq d_{(2),p} \leq \dots \leq d_{(k),p} \leq d_{(n),p}$$

Pixel which meet the stipulation in the spectral feature space, where $d_{(k),p}$ is the distance of the k-nearest neighbours and n is the number of available pixels with corresponding terrestrial data. All pixels with distances in the spectral feature space greater than $d_{(k),p}$ of the observed pixel p are ignored. With $k = 1$, only the pixel with the lowest spectral difference is been used for the further calculations. The higher the values for k that are used, the more pixels with corresponding terrestrial records affect the estimation of the target value, which contains no terrestrial records. The use of k samples means that the random scatter, caused by signal errors, may be narrowed. The distance values do only represent the differences between the spectral information of two pixels. To integrate the attribute values from the terrestrial observations, which are allocated to the k-nearest pixels, for further calculations, they have to be weighted according to their spectral distance. Hence a weighting $w_{(i),p}$ is calculated for each extracted pixel:

$$w_{(i),p} = \frac{1}{\sum_{i=1}^k \frac{1}{d_{(i),p}^2}}$$

The more similar the spectral information is, the higher is the weighting and, therefore, the influence on the attribute value, which has to be calculated.

MALTAMO & KANGAS (1998) have modified (5.5) to determine the pixel weighting

$$w'_{(t),p} = \frac{\left(\frac{1}{1 + d_{(t),p}}\right)^t}{\sum_{t=1}^k \left(\frac{1}{1 + d_{(t),p}}\right)^t}$$

where k describes the number of nearest neighbours and t influences the weighting of the distance. The bigger t is chosen, the bigger is the weighting of a pixel with a narrow spectral distance. The sum of all weightings $w'_{(t),p}$ is always 1. The attribute value of a pixel sought-after is calculated with help of the attribute values, derived from terrestrial sampling, brought into correlation with the corresponding weighted spectral data of the k-nearest pixels:

$$\hat{m}_p = \sum_{i=1}^k w_{(t),p} m_{(i),p}$$

where $m_{(i),p}$ are the terrestrially recorded values of $i = 1, \dots, k$ pixels, which are located nearest to pixel p in the spectral space. The process is repeated for every pixel and results in intensive computations, depending on the resolution of the sensor and the size of the inventory area (STÜMER, 2004).

By variation of the variables k , r , t , and a_j , influence on the estimator may be exerted. Calculations with varying values for the parameters k , r , and t are made. For an easier comparability, the value 1 was chosen for the variables a_1, a_2, \dots, a_j ." In this paper the optimal settings for the growing stock is examined and the overall accuracies of the kNN estimations were calculated.

Obtain results were evaluated on the pixel level calculating: Bias (B), Mean Square Error (MSE), Root Mean Square Error (RMSE) and procentual Root Mean Square Error (RMSE%). Bias (B) is differences between estimated and measured values:

$$B = \hat{\mu} - \mu ,$$

where $\hat{\mu}$ is an estimated value, μ the field measured value (local estimate).

The mean square error (MSE) is calculated based on variable variance $\text{Var}(\hat{\mu})$ and bias:

$$\text{MSE} = \text{Var}(\hat{\mu}) + \text{bias}^2 .$$

The root mean square error (RMSE) is determined as square root of MSE:

$$\text{RMSE} = \sqrt{\text{MSE}} .$$

The relative RMSE was calculated by using:

$$\text{RMSE}\% = \frac{\sqrt{\text{MSE}(\hat{\mu})}}{\bar{\hat{\mu}}} 100$$

where $\bar{\hat{\mu}}$ is the mean of the variable estimates.

Relative RMSE% for characteristic configurations were compared and discussed.

Beside, FRANCO-LOPEZ ET AL. (2001) found as informative to build confusion matrices for continuous attributes classifying them in attribute classes and calculate Kappa indexes as measure of agreement. A value of 0 indicates perfect agreement while a value of 0 implies that agreement is by chance. Here are analyzed confusion matrices for the growing stock classified in classes of $100 \text{ m}^3 \text{ ha}^{-1}$ and expressed Kappa values (CONGALTON, 1991). Obtained Kappa values are interpreted using following relations: < 0 - poor agreement, $0,0$ to $0,20$ - slight agreement, $0,20$ to $0,40$ - fair agreement, $0,40$ to $0,60$ - moderate agreement, $0,60$ to $0,80$ - substantial agreement and $0,80$ to $1,00$ - almost perfect agreement.

RESULTS AND DISCUSION – *Rezultati i diskusija*

The estimation efficiency depends on the natural variability and appropriate statistical methods. In the k -NN methods family it is important to examine influence of

model parameters (number of neighbors, weighting functions, other possibilities). In this paper is explored if weighted functions related to value differences and distances would affect estimation efficiency of total growing stock in high forest in Bosnia.

Firstly, values of the growing stock %RMSE depending of k nearest neighbours for different combination of value differences and distance weight are examined (Fig. 3). The typical behaviour of RMSEs is obtained resulting with sharp error decrease for the closer neighbors ($k < 3-5$), moderate for $k=5-7$ and value stabilization after 7 neighbours for all combinations. The highest errors are obtained in cases for $t=5$ while the lowest values were obtained for $r=1$ and $t=1$.




Figure 3. The effects of k for varying r and t on %RMSE
 Slika 3. Zavisnost relativne greške (%) procjene od k , r i t

The mean estimates on the pixel level are overestimated for all combination k , r and t . Procentual RMSEs were in the interval of 54 to 57%. Errors were affected by t weights resulting with higher values of biases and RMSEs for higher distance weight (Table 1).

Table 1. Evaluation results for different weight values r and t (pixel level)
 Tabela 1. Evaluacija rezultata za različite pondere r i t (nivo pixela)

r	t	Mean	Bias	SE	RMSE	RMSE%
1	1	339.79	-8.48	183.51	183.71	54.06
2	2	339.45	-8.17	183.29	183.47	54.05
2	5	343.84	-12.54	192.79	193.20	56.19
5	2	339.34	-8.04	184.16	184.34	54.32
5	5	343.20	-11.90	192.60	192.96	56.22

The RMSE-s depending of value difference weights (r) (const. $k=5$, $t=2$) and distance weights (t) (const. $k=5$, $r=2$) are presented in Figure 4.a) and 4.b). It is visible that the minimal RMSEs are obtained for $r=1$ and $t=1$ respectively. The %RMSE depending of r started with minimal value, then showed sharp increase giving importance to closer value differences with further moderate error variation (Fig. 4a). The distance weight function for constant k and r resulted with minimal value of %RMSE for t equal 1 (Fig. 4b).




Figure 4 a. %RMSE for r ($k=5, t=2$); b. %RMSE for t ($k=5, r=2$)
 Slika 4. a. Relativna greška (%) za r ($k=5, t=2$); b. Relativna greška (%) za t ($k=5, r=2$)

Obtained results are consistent with Fig. 3 where the lowest %RMSE was achieved for $r=1$ and $t=1$. FRANCO-LOPEZ ET AL. (2001) found that it was best to use no distance weights (equal weights). Consistent results related to distance weights are obtained for basal area ($r=2$) in STÜMER (2004). ÖZSAKABAŞI (2008) compared four weight functions for Euclidean distance (inverse, inverse square distance, stars and fraction) and unweighted estimates. The inverse distance based estimates resulted with the lowest error values. Findings related to t curve are published in MCROBERTS (2011) identifying the best combinations of feature variables. The author confirmed that equal neighbor weighting produced predictions comparable to more complex schemes in estimation of mean forest stem volume per unit area for small areas using a combination of forest inventory observations and Landsat Thematic Mapper (TM) imagery.

Comparison of %RMSE for different k -NN configurations employing Euclidean and Mahalanobis distances is given in Figure 5. The obtained results show consistent %RMSE behavior with the lowest values for unweighted Euclidean distance while Mahalanobis distances resulted with higher %RMSEs.




Figure 5. %RMSEs for weighted Euclidean (ED) and (modified) Mahalanobis distances (MMD, MD)

Slika 5. Relativna greška (%) za ponderisane Euklidovu (ED) i (modificiranu) Mahalanobisovu distance (MMD, MD)

ÖZSAKABAŞI (2008) compared four distances (Euclidian, Mahalanobis, diagonal Mahalanobis and Manhattan) and concluded that Euclidian gave the best results too.

Further, for different configuration using weighting functions the estimates for high forest are generated and mapped. The growing stock estimates are classified in volume classes of $100 \text{ m}^3 \text{ ha}^{-1}$ and determined Kappa value. Classified thematic map generated by different *k*-NN configurations using weighting functions and the map obtained in regular forest inventory in high forest on forest management enterprise level (FEI) were compared. Kappa values ranged from 0.53 to 0.56 with values about 0,53 for weighted Euclidian distances and 0,55 for Mahalanobis distance (Table 2).

Table 2 Kappa agreements for growing stock estimates generated using weighting functions (the high forest)

Tabela 2 Kappa pokazatelji za procjene drvne zalihe pri različitim ponderskim funkcijama (visoke šume)

Distance	Weights		Kappa
	Value difference	Distance	
	<i>r</i>	<i>t</i>	
Euclidian	1	1	0.5355
	2	2	0.5343
	1	5	0.5362
	5	1	0.5309
	5	5	0.5370
Mahalanobis			0,5513

The agreement between *k*-NN and FEI estimates could be assigned as moderate. It is interesting to notice that almost perfect agreement appeared between weighted Euclidian based estimates. Also estimates based on the Mahalanobis distance achieved agreements assigned as substantial to almost perfect compared with weighted Euclidian based estimates. Both were in moderate agreement with FEI estimates as mentioned earlier (Table 3).

Table 3 Kappa agreements between growing stock estimates on the high forest level

Tabela 3 Kappa pokazatelji između procjena drvnih zaliha na nivou visokih šuma

		Weights					MD	FEI
		t1_r1	t2_r2	t5_r1	t1_r5	t5_r5		
Weights	t1_r1	1	0.95	0.95	0.93	0.95	0.79	0.54
	t2_r2		1	0.93	0.93	0.95	0.80	0.54
	t5_r1			1	0.93	0.96	0.80	0.54
	t1_r5				1.00	0.95	0.80	0.53
	t5_r5					1.00	0.80	0.54
MD							1	0.55

CONCLUSION – *Zaključci*

Here are analysed different weighting functions related to value differences and distances in *k*-NN estimates of the growing stock on pixel and aggregated level. Determined results pointed out that weighting functions affected estimates slightly as on the pixel level so on the high forest level. The obtained differences clarified effects of weighting functions and identified possible achievements in estimation efficiency. The best results on the pixel level are obtained for unweighted value difference and distance. The results are consistent with those reported in FRANCO-LOPEZ ET AL. (2001) and MCROBERTS (2011). Some research identified better results using weighting function as ÖZSAKABAŞI (2008). It is known that the specific forest and ecological conditions influence results in particular cases. The agreement between classified estimates obtained using *k*-NN for different weighting function and estimates from regular forest inventory is assigned as moderate. The weighting functions have not showed significant influence on the classified estimates agreement level. Further possibilities for improvement are reported in RATU&KANGAS (2012) where the general mean is adjusted with the realized values of neighboring observations (measured with Euclidian distance) and weighted by correlation of the errors (or variogram) as a function of distance.

ACKNOWLEDGEMENTS – *Zahvalnice*

This paper is based on previous *k*-NN research conducted in Bosnia supported from colleagues who shared common interest and enthusiasm. Support came from Germany and Italy. We would like to thank Prof. Dr. Wolfgang Stümer who provided software for weighted functions based *k*-NN estimates. The authors gratefully acknowledge Prof. Dr. Davide Travaglini, GISDATA (Sarajevo), WALD-PROJEKT (B. Krupa) and other colleagues who supported our work unselfishly by all means.

REFERENCES - *Literatura*

- BAFFETTA, F., CORONA, P., FATTORINI L. (2011): A matching procedure to improve k-NN estimation of forest attribute maps. Original Research Article. Forest Ecology and Management, In Press, Corrected Proof, Available online 20 July 2011.
- BAFFETTA, F., FATTORINI, L., FRANCESCHI, S., CORONA, P. (2009): Design-based approach to k-nearest neighbours technique for coupling field and remotely sensed data in forest surveys. Remote Sensing of Environment, 113(3):463–475, March 2009.
- BORTZ, J. (1993): Statistik für Sozialwissenschaftler. 4. Auflage. Springer, Berlin.
- CHIRICI, G. (2001): Developing an experimental survey protocol based on spatialized indicators to support forest management planning. Ph.D. thesis, Dipartimento di Scienze dell'Ambiente Forestale e delle sue Risorse. Università degli Studi della Toscana. Italy.
- CHIRICI, G., CORONA, P., MARCHETTI, M., MASELLI, F., BOTTAI, L. (2002): Spatial distribution modelling of forest attributes coupling remotely sensed imagery and GIS techniques. Reality, models and parameter estimation – the forestry scenario 2-5 June 2002, Sesimbra, Portugal.
- CONGLALTON, R. (1991): A review of assessing the accuracy of remotely sensed data. Remote Sensing of Environment 37: 35-46.
- ČABARAVDIĆ, A. (2007): Efficient Estimation of Forest Attributes with k NN. Ph.D. thesis, Faculty of Forest and Environmental Sciences, Albert-Ludwigs-Universität Freiburg im Breisgau, Germany.
- FINLEY A., MCROBERTS, R.E. (2008): Efficient k-nearest neighbour searches for multi-source forest attribute mapping. Remote Sensing of Environment, 112(5):2203–2211.
- FINLEY A., MCROBERTS, R.E., EK, A.R. (2006): Applying an efficient k Nearest neighbor search to forest attribute imputation. Forest Science, Volume 52, Number 2, April 2006, pp. 130-135(6), 52(2):130–135.
- FRANCO-LOPEZ, H., EK, A. R., BAUER, M. E. (2001): Estimation and mapping of forest stand density, volume, and cover type using the k nearest neighbors method. Remote Sensing of Environment, 77(3):251–274.
- GALLAUN, H., ZANCHI, G., NABUURS, G.J., HENGEVELD, G., SCHARDT, M., VERKERK P.J. (2010): EU-wide maps of growing stock and above-ground biomass in forests based on remote sensing and field measurements Original Research Article. Forest Ecology and Management, Volume 260, Issue 3, 30 June 2010, Pages 252-261.
- HÄRKÖNEN, S., LEHTONEN, A., EERIKÄINEN, K., PELTONIEMI, M., MÄKELÄ A. (2011): Estimating forest carbon fluxes for large regions based on process-based modelling, NFI data and Landsat satellite images Original Research Article Forest

Ecology and Management, Volume 262, Issue 12, 15 December 2011, Pages 2364-2377.

- KANGAS A., MALTAMO M. (2006): Forest inventory – Methodology and Application. Springer.
- KATILA, M. AND TOMPPPO, E. (2001): Selecting estimation parameters for the Finnish multisource National Forest Inventory. Remote Sensing of Environment, 76: 16-32.
- KÖHL M., STÜMER, W., KENTER, B., RIEDEL T. (2008): Effect of the estimation of forest management and decay of dead woody material on the reliability of carbon stock and carbon stock changes - A simulation study Original Research Article Forest Ecology and Management, Volume 256, Issue 3, Pages 229-236.
- KUTZER CH. (2007): Potential of the kNN Method for Estimation and Monitoring off-Reserve Forest Resources in Ghana. Ph.D. thesis, Faculty of Forest and Environmental Sciences, Albert-Ludwigs-Universität Freiburg im Breisgau, Germany.
- KOUKAL, T. (2004): Nonparametric Assessment of Forest Attributes by Combination of Field Data of the Austrian Forest Inventory and Remote Sensing Data. Ph.D. thesis, Universität für Bodenkultur Wien.
- KOHL, M.; STUMER, W.; KENTER, B.; RIEDEL, T. (2008): Effect of the estimation of forest management and decay of dead woody material on the reliability of carbon stock and carbon stock changes—A simulation study. Forest Ecology and Management. 256: 229-236.
- MALTAMO, M. & KANGAS, A., (1998): Methods based on k-nearest neighbour regression in the prediction of basal area diameter distribution. Can. J. For. Res. 28: 1107-1115.
- MASELLI, F., CHIRICI, G., BOTTAI, L., CORONA, P., MARCHETTI, M. (2005): Estimation of mediterranean forest attributes by the application of k-NN procedures to multitemporal landsat ETM+ images. International Journal of Remote Sensing, 26(17):3781–3796.
- MCRROBERTS, R., E. TOMPPPO, A. FINLEY, HEIKKINEN J. (2007): Estimating areal means and variances of forest attributes using the k-Nearest neighbors technique and satellite imagery. Remote Sensing of Environment, 111(4):466–480.
- MCRROBERTS. R.E. (2008): Using satellite imagery and the k-nearest neighbors technique as a bridge between strategic and management forest inventories. Remote Sensing of Environment, 112(5):2212–2221, May 2008.
- MCRROBERTS, R.E. (2011): Estimating forest attribute parameters for small areas using nearest neighbors techniques Original Research Article Forest Ecology and Management, In Press, Corrected Proof, Available online 3 September 2011.
- MCRROBERTS, R.E., MAGNUSSEN, S., TOMPPPO, E.O., CHIRICI G. (2011): Parametric, bootstrap, and jackknife variance estimators for the k-Nearest Neighbors

- technique with illustrations using forest inventory and satellite image data Original Research Article Remote Sensing of Environment, In Press, Corrected Proof, Available online 27 August 2011.
- NABUURS, G.J., HENGEVELD, G.M., VAN DER WERF, D.C., HEIDEMA A.H. (2010): European forest carbon balance assessed with inventory based methods—An introduction to a special section Original Research Article. Forest Ecology and Management, Volume 260, Issue 3, 30 June 2010, Pages 239-240.
- ÖZSAKABAŞI, F. (2008): Classification of forest areas by k nearest neighbor method: case study, Antalya. Master thesis. Geodetic and geographic information technologies department, Middle East Technical University.
- RÄTY, M., KANGAS A. (2012): Comparison of k-MSN and kriging in local prediction Original Research Article Forest Ecology and Management, Volume 263, 1 January 2012, Pages 47-56.
- STÜMER, W., KENTER, B., KÖHL M. (2010): Spatial interpolation of in situ data by self-organizing map algorithms (neural networks) for the assessment of carbon stocks in European forests. Original Research Article. Forest Ecology and Management, Volume 260, Issue 3, 30 June 2010, Pages 287-29.
- STÜMER, W. (2004): Kombination von terrestrischen Aufnahmen und Fernerkundungsdaten mit Hilfe der kNN-Methode zur Klassifizierung und Kartierung von Wäldern. PHD. Fakultät für Forst-, Geo- und Hydrowissenschaften der Technischen Universität Dresden.
- TOMPPONEN, E., KORHONEN, K.T., HEIKKINEN, J. AND YLI-KOJOLA, H. (2001): Multisource inventory of the forests of the Hebei Forestry Bureau, Heilongjiang, China. *Silva Fennica* 35: 309-328.
- TOMPPONEN, E. & HALME, M. (2004): Using coarse scale forest variables as ancillary information and weighting of variables in k-NN estimation: a genetic algorithm approach. *Remote Sensing of Environment* 92: 1-20.
- TOMPPONEN, E., OLSSON, H., STAHL, G., NILSSON, M., HAGNER, O., KATILA, M. (2008): Combining national forest inventory field plots and remote sensing data for forest databases. *Remote Sensing of Environment*, 112(5):1982–1999.

SAŽETAK

Kombinovane inventure šuma u kojima se koriste daljinska istraživanja omogućile su podizanja efikasnosti procjena šumskih resursa primjenom različitih metodoloških pristupa. Najčešće korišteni metod procjene najvažnijih šumskih atributa je neparametrijski metod k najbližih susjeda koji je pokazao visoku efikasnost pri integraciji terenskih snimanja i LANDSAT satelitskih snimaka. Metod pruža mogućnost prilagođavanja konkretnim prilikama tj. podizanju efikasnosti procjena pomoću ponderiranih funkcija koje daju različit značaj vrijednostima terenskih snimanja. U ovom radu analizirane su ponderirane funkcije s obzirom na razlike vrijednosti koje se javljaju u cijelom uzorku i udaljenost registrovanih vrijednosti u multidimenzionalnom prostoru spektralnih kanala. Analizirane su greške procjena drvene zalihe na nivou pixela i na nivou visokih šuma kao kategorije. Ustanovljeno je da ponderisanje razlika vrijednosti i udaljenosti nije rezultiralo povećanom efikasnošću procjena na nivou pixela. Neparametrijske procjene drvnih zaliha u visokim šumama dostigle su umjeren stepen slaganja sa procjenama dobijenim redovnom taksacijom šuma kako za neponderisane tako i za ponderisane procjene. Između neponderisanih i ponderisanih procjena javlja se gotovo potpuno slaganje. Dobijeni rezultati podržavaju jednostavnije procedure procjene i potvrđuju predhodno izvedena rješenja. Dalja istraživanja treba usmjeriti na napredne metodološke mogućnosti i njihovo prilagođavanje za korištenje u konkretnim uvjetima.